USDA United States Department of Agriculture | Forest Service | Northeastern Area State and Private Forestry | 180 Canfield Street Morgantown, WV 26505-3101

369*File Code: 3400
Date: October 21, 2008

Mr. Richard Patterson, Director
Northeastern Area State and Private Forestry
Grey Towers National Historic Landmark
ATTN: Elizabeth Hawke, Horticulturist
151 Grey Towers Drive
P.O. Box 188
Milford, PA 18337-0188

Dear Mr. Patterson:

Enclosed is the gypsy moth biological evaluation for Grey Towers National Historic Landmark.

In brief, gypsy moth populations are sufficient to cause light defoliation on 87 acres. We are recommending a single application of *Bacillus thuringienis* variety *kurstaki (Btk)*. With good timing and proper application, gypsy moth defoliation should be prevented at Grey Towers in 2009.

Please contact me at 304-285-1555 if you have any questions regarding this gypsy moth biological evaluation.

Sincerely,

Rodney L. Whiteman

RODNEY L. WHITEMAN
Forester
Forest Health Protection

Enclosure

Cc: Tim Marasco, PA BOF
Robert Lueckel, MFO

BIOLOGICAL EVALUATION OF GYPSY MOTH

At

GREY TOWERS NATIONAL HISTORIC LANDMARK

2008

Prepared by

Rodney L. Whiteman
Forester

USDA Forest Service
Forest Health Protection
Morgantown, WV 26505

October 2008

ABSTRACT

On September 25, 2008, USDA Forest Service personnel conducted a gypsy moth egg mass survey at Grey Towers National Historic Landmark. The purposes of the survey were to determine gypsy moth population densities, evaluate the efficacy of this year's treatments, assess the potential for defoliation and the need for treatment in 2009. Current populations are sufficient to cause light defoliation on 87 acres. Treatment is recommended to prevent defoliation, possible branch dieback and possible tree mortality.

METHODS

Gypsy moth survey plots were randomly selected based upon available host trees (oak species), size of sample area and uniformity between egg mass counts. At each sample point, a 1/40th acre fixed radius plot was established. The plots consisted of a tally of all the new (2008) egg masses observed on the overstory trees, understory vegetation, ground litter and duff. The total number of egg masses observed for each plot was multiplied by 40 to determine the number of egg masses per acre. Egg mass lengths were measured at the plots to determine the overall "health" of the existing population and as a measure of egg mass fecundity.

RESULTS

The location of the survey plots along with the 2008 *Bacillus thuringienis* variety *kurstaki (Btk)* treatment area are shown in Figure 1. The summarized results of the survey are presented in Table 1. In brief, egg mass densities ranged from 40-3400 and averaged 1273 egg masses per acre. Overall egg mass lengths tended to be moderate in size, ranging from 15-42 mm and averaging 25 mm.

Egg mass densities at Grey Towers have actually increased 18 percent from the pre-treatment level (2007) of 1080 to their current level. However, egg mass lengths were much longer in 2007 when they averaged 34 mm.

DISCUSSION

The basic guidelines used to evaluate the risk of defoliation include: previous defoliation events; number of egg masses/acre; size and condition of the egg masses; available preferred food; and risk of larval blow-in following egg hatch. Potential defoliation is categorized as; light (30-50 percent); and heavy (51-100 percent). Defoliation less than 30 percent generally has little impact on trees and can not be detected through aerial surveys.

The survey results indicate that light defoliation is likely on 87 acres (Figure 2).
This conclusion is further supported when egg density is used as a means of predicting defoliation. Moore and Jones (1987) found that estimating the mean fecundity would increase the precision of gypsy moth density estimates and that a linear relationship exists

between egg mass length and fecundity. Further work by Liebhold et al., (1993) demonstrates that the product of the mean egg mass length (in mm) and egg mass density provides a more precise means of estimating population densities and prediction defoliation. Using Liebhold's model, Figure 3 shows how this information can be used to correlate the predicted defoliation of an area. Accordingly, the estimated egg mass density of 1273 egg masses per acre (average egg mass density) x 25 mm (average egg mass length) translates to a projected defoliation level of about 36 percent (light defoliation). Because egg mass densities and the host type are not evenly distributed, actual defoliation will vary from tree to tree but will be predominately light throughout Grey Towers.

Based on existing egg mass densities and the general size of egg masses, gypsy moth populations appear to be fairly static throughout Grey Towers. The average egg mass length is 25 mm. Egg masses larger than 25 mm typically indicate healthy populations with no obvious stress from either the gypsy moth nucleopolyhedrosis virus (NPV) or the *Entomophaga maimaiga* fungus, two of the primary natural control agents that often express themselves in declining or stressed populations. There was no evidence that either one of these entomopathogens had significant impacts at Grey Towers in 2008. Although it is still possible that either the gypsy moth fungus or the NPV could cause the general collapse of the gypsy moth population next year, it is unlikely that populations will collapse prior to a significant defoliation event occurring in 2009.

Predicting the extent of tree mortality that would occur after one year's defoliation is difficult, however, a stand of trees that is not stressed by other agents during or immediately following a single heavy defoliation will likely pull through with only minor branch dieback and minimal mortality. Trees that are defoliated in excess of 60 percent normally refoliate the same growing season. Such events cause the trees to expend valuable energy reserves to refoliate, and consequently cause the trees' health to deteriorate. Depending on the condition of the trees at the time of defoliation, reduced growth, mast abortion, branch dieback or in some cases tree mortality, has occurred following a single year of heavy defoliation. Should subsequent defoliation occur the following year, the impact is compounded. Trees that receive light defoliation (<50 percent) are not likely to refoliate and there is probably no significant impact other than a reduction in growth, reduction of mast and possibly some minor branch dieback.

Trees at greater risk are those that are presently stressed from other factors, such as soil compaction from roads, sidewalks, parking lots, machinery and/or heavy foot travel; over maturity; drought; shock due to recent timber cutting activities; previous year(s) defoliation; and other insect and disease related problems. Grey Towers experienced a prolonged and severe drought during the 2007 growing season, and again late in the 2008 growing season. Less than 1 acre of the defoliation was detected at Grey Towers in both 2007 and 2008.

The Allegheny National Forest (1988) and the West Virginia Division of Forestry (1997) provide examples of the potential tree mortality that can occur. On the Allegheny National Forest, untreated stands consisting of 40-80 percent oak, the average loss of basal area (mainly oaks) was about 16 percent (range 3-28 percent) following one year of defoliation and 26 percent (range 10-43 percent) after two consecutive years of defoliation. In a 1986

study area in eastern West Virginia where oak species accounted for 63-78 percent of the species composition, a loss of 25 percent of the total oak saw timber and 14 percent of the total oak pole timber occurred after one year of light to heavy defoliation. In these examples, droughty conditions likely contributed to the level of mortality.

Based on observations of the existing health of the forested areas at Grey Towers and the factors mentioned above, small scattered areas of tree mortality may occur if defoliation occurs. Mortality will be more likely if adequate rainfall is not received during the 2009 growing season and or defoliation occurs on trees that were previously defoliated.

Management Options

In 2009, three management options have been evaluated for managing gypsy moth populations at Grey Towers. The intervention options are offered based upon the following two treatment objectives: 1) protect host tree foliage to prevent branch dieback and tree mortality; and 2) reduce gypsy moth population below the treatment threshold. Each is discussed below.

No Action Option

It is possible that gypsy moth populations could collapse on their own due to the presence of nucleopolyhedrosis virus (NPV) or the more recently recognized fungal pathogen, *Entomophaga maimaiga*. In areas with defoliating levels of gypsy moth populations, viral epizooics generally manifest themselves after significant tree defoliation has already occurred. Gypsy moth populations will usually peak in 2-3 years once they reach levels and then collapse as a result of NPV or fungal activity. Residual populations following such a collapse will likely remain at low densities for 3-6 years before rebuilding to defoliating levels.

Although it is not possible to accurately assess such events with the defoliating levels and then collapse as a result of NPV or fungal activity. Residual populations information at hand, it is unlikely that a collapse will occur in 2009 since the egg masses appear to be fairly healthy.

Large numbers of gypsy moth caterpillars and defoliation has been shown to impact competing native herbivore arthropods. Sample et al., (1996) showed short-term impacts of both species richness and abundance occurred following light to moderate defoliation events in study plots in West Virginia. It is likely that impacts would be greater as the size of the area and intensity of defoliation increases and be more long term, should extensive tree mortality occur.

Should this option be selected, it is likely that light defoliation will occur at Grey Towers in 2009 (Figure 2).

Microbial Insecticide Option

***Btk*:** The only biological insecticide currently registered and commercially available for gypsy moth control is the microbial insecticide *Bacillus thuringienis* variety *kurstaki (Btk)*. This insecticide is available through several manufacturers and has been used extensively in suppression projects throughout the U.S. in both forested and residential areas. *Btk* is a bacterium that acts specifically against lepidopterous larvae as a stomach poison and therefore must be ingested. The major mode of action is by mid-gut paralysis which occurs soon after feeding. This results in a cessation of feeding, and death by starvation. *Btk* is persistent on foliage for about 7-10 days.

Btk has been shown to impact other non-target caterpillars that are actively feeding at the time of treatment. An example of the potential impacts is provided by a study conducted by Miller (1990) in Oregon and Samples, et al., (1996) in West Virginia. Miller's study involved a large scale (5,000 acres) eradication program where three consecutive applications of *Btk* were applied within a single season. On Garry oak, Miller found that species richness was significantly reduced in treated areas during all 3 years of the study while the total number of immature native Lepidoptera rebounded after the second year. In the Sample study, the areas treated with *Btk* were 50 acre plots and only a single treatment applied. Here too, both species richness and the total numbers of native macro-lepidopterous caterpillars and adults were reduced but only for less than 1 year. The difference in duration of the impacts between these studies is probably the result of the number of treatment applications applied and the size of the treatment area involved.

Btk formulations are available as flowable concentrates, wettable powders, and emulsifiable suspensions. The normal application rates range from 24-36 billion international units (BIUs) per acre in a single or double application. *Btk* can be applied either undiluted or mixed with water for a total volume of ½ -1 gallon per acre. With proper application, foliage protection and some degree of population reduction can be expected with one application and with two applications both foliage protection and a greater degree of population reduction are likely.

Because *Btk* is a biological insecticide, the degree of population reduction varies and may depend on, at least in part, the selected application rate, relative health of the population (building vs. declining), population densities, weather (rain and temperature), the feeding activity of the larvae following treatment, and the actual potency of the product.

Gypchek: A second microbial insecticide that is registered and available in limited quantities is the formulated nucleopolyhedrosis virus called Gypchek. This product is not available commercially but is produced in limited quantities by a cooperative effort of the USDA Forest Service and the Animal Plant Health Inspection Service (APHIS). The active ingredient in Gypchek formulations has a very narrow host range (lymnatriids) and occurs naturally in gypsy moth populations. Normally the virus reaches epizootic proportions when gypsy moth populations reach high densities as a result of increased transmission within and

between gypsy moth generations. The application of Gypchek to gypsy moth populations simply expedites this process by increasing the exposure of the virus at an earlier stage. Healthy, feeding gypsy moth caterpillars become infected by ingesting contaminated foliage and soon stop feeding and die.

The efficacy of Gypchek treatments to reduce gypsy moth populations has been quite variable. Because of the short period of viral activity on foliage (3-5 days) as well as other biological factors such as feeding activity and weather conditions, it has been difficult at best to project treatment efficacy. Most often foliage protection can be achieved but significant reductions in gypsy moth densities do not always occur. Should inadequate population reduction occur, areas would need to be treated again the following year.

The normal application rate of Gypchek is 4×10^{11} occlusion bodies (OB's) per acre applied in a single application or 2×10^{11} OB's per acre applied in a double application. Due to the limited supply, priority is first given to state and federal cooperators that need to deal with federally listed threatened and endangered species associated with gypsy moth treatments. There are, however, sufficient quantities of Gypchek currently available for 2009 should this insecticide be preferred for use at Grey Towers.

Chemical Insecticide Option

The third option is to us a chemical insecticide to control gypsy moth populations. There is currently one chemical insecticide registered for control of gypsy moth populations and approved by the USDA Forest Service for use in cooperative gypsy moth control programs.

Dimilin® (diflubenzuron) is the most widely used chemical insecticide in gypsy moth suppression projects in the U.S. and is registered by the EPA for use in residential areas. Diflubenzuron (DFB) is an insect growth regulator that disrupts the normal molting processes of the larvae. The mode of action is to inhibit the formation chitin, a necessary component of the outer cuticle which causes the affected larvae to die during the molt following treatment. The methods of uptake is primarily by ingestion, however some research has indicated the possibility of absorption through the cuticle as well. DFB is relatively persistent on foliage (24 days) which increases the efficacy on gypsy moth populations but also exposes non-target insects, particularly caterpillars, for a greater period of time.

DFB is extremely toxic to some aquatic invertebrates and the label prohibits the application over open water or wetlands. DFB is available as an oil based liquid formulation (Dimilin® 4L) and is normally applied in a single application at the standard rate of 1-2 ounces of formulated material per acre. With proper application, foliage protection and a significant population reduction can be expected. The need for treatment of residual populations the following year is normally not necessary.

Alternatives

With the previously described options in mind, the following alternatives are offered:

Alternative 1. - No action.

Alternative 2. - One aerial application of *Btk* at the rate of 36 BIUs in a total mix of ¾ gallon per acre.

Alternative 3. - Two aerial applications of *Btk*, as in alternative 2, applied 4-7 days apart.

Alternative 4. - One aerial application of Gypchek at the rate of 4×10^{11} OB's in a total mix of ¾ gallon per acre.

Alternative 5. - Two aerial applications of Gypchek at the rate of 2×10^{11} OB's in a total mix of ¾ gallon per acre, applied 3-5 days apart.

RECOMMENDATIONS

As previously stated, gypsy moth populations at Grey Towers are sufficient to cause 87 acres of defoliation in 2009. To protect tree foliage, and prevent possible branch dieback and prevent tree mortality, our recommendation is alternative 2 (a single application of Btk).

This recommendation is based on the following considerations:

1) A single application of Btk is likely to provide both better foliage protection and a significant gypsy moth population reduction.

2) Current gypsy moth populations don't warrant a double application of Btk (alternative 3) or a single application of DFB (Alternative 6).

3) A single application of Btk is more likely to provide both foliage protection and a significant gypsy moth population reduction than a single application of Gypchek (Alternative 4).

4) A single application of Btk is more economical choice than Gypchek.

REFERENCES

Allegheny National Forest, Warren, PA. 1998. Gypsy moth caused oak mortality – Allegheny National Forest, 1988. USDA Forest Service internal report prepared by Forest Pest Management staff, Morgantown, WV. Unp.

Gottschalk, K.W. 1990. Gypsy moth impacts on mast production, *In*: McGee, Charles E. Ed. Proceedings of the Workshop, southern Appalachian Mast Management; 1989 August 14-16; Knoxville TN; University of Tennessee; 42-50.

Liebhold, A.M., Simons, E.E., Sior, A., and Unger, J.D. 1993. Forecasting defoliation caused by the gypsy moth from field measurements. Environ. Entomol. 22(1): 26-32.

Miller, J.C. 1990. Field assessment of the effects of a microbial pest control agent on non-target Lepidoptera. American Entomologist 36:2, 135-139.

Moore, K.E.B. and Jones, C.G. 1987. Field estimation of fecundity of gypsy moth (Lepidoptera: Lymnatriidae). Environ, Entomol. 16: 165-167.

Sample, B.E., Butler, L., Zivkovich, C., Whitmore, R.C., and Reardon, R.C. 1996. Effects of *Bacillus thuringiensis* Berliner var. *Kurstaki* and defoliation by gypsy moth [*Lymantria dispar* (L.) (Lepidoptera: lymnatriidae)] on native arthropods in West Virginia. The Canadian Entomologist 128:573-592.

West Virginia Division of Forestry. 1997. *In* 1997 Cooperative State-County-Landowner Gypsy Moth Suppression Program in West Virginia. 3p. (Brochure).

Table 1. – Results of the gypsy moth egg mass survey conducted at Grey Towers on September 25, 2008.

Plot #	#em/acre	em size (mm)
1	120	32
2	480	14, 20, 35
3	40	--
4	920	15, 29
5	3400	20, 25, 42
6	2680	15, 24, 30

em/acre range = 40-3400
em/acre average = 1273

em size range (mm) = 15-42
em size average (mm) = 25

Figure 1. -- Location of the gypsy moth survey plots established on September 25, 2008, at Grey Towers National Historic Landmark, along with the 2008 treatment area.



Figure 2. -- Area where defoliation is likely/recommended treatment area at Grey Towers National Historic Landmark.



Figure 3.—Predicted defoliation at Grey Towers National Historic Landmark in 2009.



Scatter plot of the product of mean egg mass length and egg mass density versus mean defoliation. Extracted from Liebhold et al. (1993).